Plant Pathology (1995) **44**, 1085–1092

Author index

Agapov V. *see* Reuveni M.
Aguirreolea J., Irigoyen J., Sánchez-D'az M. & Salaverri J. Physiological alterations in pepper during wilt induced by *Phytophthora capsici* and soil water deficit, 587
Ahmed H.U., Mundt C.C. & Coakley S.M. Host-pathogen relationship of geographically diverse isolates of *Septoria tritici* and wheat cultivars, 838
Airede C. *see* Mepsted R.
Alcázar M.D. *see* Candela M.E.
Al-Kaff N.S. & Covey S.N. Biological diversity of cauliflower mosaic virus isolates expressed in two Brassica species, 516
Al-Kherb S.M., Fininsa C., Shattock R.C. & Shaw D.S. The inheritance of virulence of *Phytophthora infestans* to potato, 552
Almela L. *see* Candela M.E.
Ammar E.D., Gingery R.E. & Madden L.V. Transmission efficiency of three isolates of maize stripe tenuivirus in relation to virus titre in the planthopper vector, 239
Amoah B.K., Rezanoor H.N., Nicholson P. & MacDonald M.V. Variation in the *Fusarium* section *Liseola*: pathogenicity and genetic studies of isolates of *Fusarium moniliforme* Sheldon from different hosts in Ghana, 563
Andrivon D. *see* Newton A.C.
Antoniou P.P., Tjamos E.C. & Panagopoulos C.G. Use of soil solarization for controlling bacterial canker of tomato in plastic houses in Greece, 438
Arancibia R.A., Valverde R.A. & Can F. Properties of a cryptic virus from pepper (*Capsicum annuum*), 164
Archer S.A. *see* McQuilken M.P.
Arias R. *see* Maddison A.C.
Arnold D.L. *see* Claxton J.R.
Asher M.J.C. *see* Barr K.J.

Balali G.R., Neate S.M., Scott E.S., Whisson D.L. & Wicks T.J. Anastomosis group and pathogenicity of isolates of *Rhizoctonia solani* from potato crops in South Australia, 1050
Bandyopadhyay R. *see* Musabyimana T.
Bannister P. *see* Eason J.R.
Barbara D.J. *see* Morton A.
Barr K.J., Asher M.J.C. & Lewis B.G. Resistance to *Polymyxa betae* in wild *Beta* species, 301
Barton J.E. *see* Yang H.A.
Bejarano-Alcázar J. *see* Melero-Vara J.M.
Berggren G.T. *see* Kousik C.S.
Berio T. *see* Migheli Q.
Bertolini P. *see* Tian S.P.
Bevan J.R., Taylor J.D., Crute I.R., Hunter P.J. & Vivian A. Genetics of specific resistance in pea (*Pisum sativum*) cultivars to seven races of *Pseudomonas syringae* pv. *pisi*, 98
Bharathan N. *see* Ramos L.J.
Billard L. *see* Chakraborty S.
Blakesley D. *see* Claxton J.R.
Blanco-Lùpez M.A. *see* Melero-Vara J.M.
Boscia D., Elicio V., Savino V. & Martelli G.P. Production of monoclonal antibodies to grapevine fleck virus, 160
Boudon-Padieu E. *see* Seddas A.
Brasier C.M., Rose J. & Gibbs J.N. An unusual *Phytophthora* associated with widespread alder mortality in Britain, 999
Broglio E.P. *see* Satyanarayana T.
Brophy L.S. *see* Mundt C.C.
Brown A.E. *see* Fletcher J.T.
Brown J.K.M. Recombination and selection in populations of plant pathogens, 279
Brown J.K.M. & Jessop A.C. Genetics of avirulences in *Erysiphe graminis* f.sp. *hordei*, 1039
Brunet Y. *see* Rott P.
Budge S.P. *see* McQuilken M.P.
Burdon J.J. & Roberts J.K. The population genetic structure of the rust fungus *Melampsora lini* as revealed by pathogenicity, isozyme and RFLP markers, 270
Butler D.R., Reddy R.K. & Wadia K.D.R. Single-plant chambers to control humidity for pathological studies, 1
Butler D.R., Wadia K.D.R. & Reddy R.K. Effects of humidity, leaf wetness, temperature and light on conidial production by *Phaeoisariopsis personata* on groundnut, 662
Butt D.J. *see* Xu X.-M.

Café Filho A.C. & Duniway J.M. Dispersal of *Phytophthora capsici* and *P. parasitica* in furrow-irrigated rows of bell pepper, tomato and squash, 1025
Can F. *see* Arancibia R.A.
Candela M.E., Alcázar M.D., Esp'n A., Egea C. & Almela L. Soluble phenolic acids in *Capsicum annuum* stems infected with *Phytophthora capsici*, 116
Caponero A., Contesini A.M. & Iacobellis N.S. Population diversity of *Pseudomonas syringae* subsp. *savastanoi* on olive and oleander, 848
Carder J.H. *see* Morton A.
Cassells A.C. & Walsh M. Screening for *Sclerotinia* resistance in *Helianthus tuberosus* L. (Jerusalem artichoke) varieties, lines and somaclones, in the field and *in vitro*, 428
Castillo P., Jimenez-Diaz R.M., Gomez-Barcina A. & Vovlas N. Parasitism of the root-lesion nematode *Pratylenchus thornei* on chickpea, 728
Castoria R., Fanelli C., Zoina A. & Scala F. Analysis of fatty acids in lipids of *Verticillium dahliae* and induction of lubimin accumulation in eggplant, 791
Ceniza M.S. *see* Sugimura Y.
Cervantes L.A. *see* Spotts R.A.
Chakraborty S. & Billard L. Quantitative relationships between *Colletotrichum gloeosporioides* infection of *Stylosanthes scabra* and weather under field conditions, 63
Chalutz E. *see* Wisniewski M.
Chartier R. *see* Keck M.
Clarkson J.M. *see* Claxton J.R.
Claxton J.R., Arnold D.L., Blakesley D. & Clarkson J.M. The effects of temperature on zoospores of the crook root fungus *Spongospora subterranea* f.sp. *nasturtii*, 765
Clifford B.C. *see* Roderick H.W.
Coakley S.M. *see* Ahmed H.U.
Cockerell V., Rennie W.J. & Jacks M. Incidence and control of barley leaf stripe (*Pyrenophora graminea*) in Scottish barley during the period 1987-1992, 655
Cole D.L. *see* Edwards C.
Contesini A.M. *see* Caponero A.
Cook R.J. New honorary member of the British Society for Plant Pathology, 405
Cooper R.M. *see* Mepsted R., Resende M.L.V.
Covey S.N. *see* Al-Kaff N.S.
Cowley T. *see* Walters D.R.
Cowling W.A. *see* Willamson P.M.
Cristinzio G., Rotino G.L. & Scala F. Investigation of *in-vitro* methods for measuring eggplant resistance to *Verticillium dahliae*, 704
Crute I.R. *see* Bevan J.R.
Cullum F.J. *see* Simpson M.J.A.

Dai Qun & Sun Zhi-Huan Suppressive effects of N-triacontanol on symptoms of mulberry dwarf disease and on the causal phytoplasma, 979
Dann E.K. & Deverall B.J. Effectiveness of systemic resistance in bean against foliar and soilborne pathogens as induced by biological and chemical means, 458
Day J.P. & MacDonald M.V. Plant-pathogen interactions of sunflower and *Macrophomina phaseolina in vitro* and *in vivo*, 261
de Cal A., Pascual S., Larena I. & Melgarejo P. Biological control of *Fusarium oxysporum* f.sp. *lycopersici*, 909
de Vallavieille-Pope C. *see* Lannou C.
Deacon J.W. *see* Jones E.E.
Défago G. *see* Maurhofer M.
Dempster L.C. & Holmes S.J.I. The incidence of strains of barley yellow dwarf virus in perennial ryegrass crops in south-west and central Scotland, 710
Deverall B.J. *see* Dann E.K.
Diourte M., Starr J.L., Jeger M.J., Stack J.P. & Rosenow D.T. Charcoal rot (*Macrophomina phaseolina*) resistance and the effects of water stress on disease development in sorghum, 196
Dixon G.R. *see* Kenyon D.M.
Drenth A., Janssen E.M. & Govers F. Formation and survival of oospores of *Phytophthora infestans* under natural conditions, 86
Drepper W.J. *see* Hau B.
Droby S. *see* Wisniewski M.
Duncan J.M. & Kennedy D.M. Effect of temperature and host genotype on the production of inoculum by *Phytophthora fragiariae* var. *fragiariae* from the roots of infected strawberry plants, 10
Duncan R.A., Stapleton J.J. & Leavitt G.M. Population dynamics of epiphytic mycoflora and occurrence of bunch rots of wine grapes as influenced by leaf removal, 956
Duniway J.M. *see* Café Filho A.C.
Dyer P.S. & Lucas J.A. Incidence of apothecia of *Tapesia yallundae* at set-aside sites in England and sensitivity of the ascospore offspring to the fungicides benomyl and prochloraz, 796

Eason J.R., Jameson P.E. & Bannister P. Virulence assessment of *Rodococcus fascians* strains on pea cultivars, 141
Edgar J.A. *see* Willamson P.M.
Edwards C., Strange R.N. & Cole D.L. Accumulation of isoflavonoid phytoalexins in leaves of *Arachis hypogaea* differing in reaction to rust (*Puccinia arachidis*) and early leafspot (*Cercospora arachidicola*), 573
Egea C. *see* Candela M.E.
Eguchi S. *see* Sugimura Y.
Eilam Y. *see* Wisniewski M.
Elena K. & Tjamos E.C. Vegetative compatibility groups of *Fusarium oxysporum* f.sp. *dianthi* from plants and the rhizosphere of carnation in Greece, 148
Elicio V. *see* Boscia D.
Erwin D.C. *see* Howell A.B.
Esp'n A. *see* Candela M.E.

Fanelli C. *see* Castoria R.
Feldmann P. *see* Rott P.
Fenlon J.S. *see* McQuilken M.P.
Fiddaman P.J. & Rossall S. Selection of bacterial antagonists for the biological control of *Rhizoctonia solani* in oilseed rape (*Brassica napus*), 695
Figueroa L., Fitt B.D.L., Shaw M.W., McCartney H.A. & Welham S.J. Effects of temperature on the development of light leaf spot (*Pyrenopeziza brassicae*) on oilseed rape (*Brassica napus*), 51
Figueroa L., Fitt B.D.L., Welham S.J., Shaw M.W. & McCartney H.A. Early development of light leaf spot (*Pyrenopeziza brassicae*) on winter oilseed rape (*Brassica napus*) in relation to temperature and leaf wetness, 641
Fininsa C. *see* Al-Kherb S.M.
Fitt B.D.L. *see* Figueroa L.
Fletcher J.T., Jaffe B., Muthumeenakashi S., Brown A.E. & Wright D.M. Variations in isolates of *Mycogone perniciosa* and in disease symptoms in *Agaricus bisporus*, 130
Flood J. *see* Mepsted R., Resende M.L.V.

Garcia Heriz M. *see* Geyl L.
Garibaldi A. *see* Migheli Q.
Gaunt R.E. *see* Sanoamuang N.
Gelie B. *see* Jamaux I.
Geyl L., Garcia Heriz M., Valentin P., Hehn A. & Merdinoglu D. Identification and characterization of resistance to rhizomania in an ecotype of *Beta vulgaris* subsp. *maritima*, 819
Gibbs J.N. *see* Brasier C.M.
Gingery R.E. *see* Ammar E.D.
Gomez-Barcina A. *see* Castillo P.
Goodwin P.H., Hsiang T., Xue B.G. & Liu H.W. Differentiation of *Gaeumannomyces graminis* from other turf-grass fungi by amplification with primers from ribosomal internal transcribed spacers, 384
Gorris L.G.M. *see* Smid E.J.
Govers F. *see* Drenth A.
Goyeau H. *see* Lannou C.
Gullino M.L. *see* Migheli Q.
Gulya T.J. *see* Virányi F.
Gustafsson M. *see* Olbe M.

Haas D. *see* Maurhofer M.
Habtu Assefa, van den Bosch F., Zadoks J.C. Focus expansion of bean rust in cultivar mixtures, 503
Hale C.N. *see* Kearns L.P.
Harrison J.G. *see* Wallace A.
Harville B.G. *see* Kousik C.S.
Hau B., Drepper W.J., Prasertkit O., Kranz J. & Renfro B.L. Temporal and spatial aspects of the epidemiology of sorghum downy mildew on maize, 897
Havis N.D. *see* Walters D.R.
Haware M.P. *see* Stevenson P.C.
Hehn A. *see* Geyl L.
Heiberg N. Control of root rot of red raspberries caused by *Phytophthora fragariae* var. *rubi*, 153
Helfer S. *see* Kenyon D.M.
Hijwegen T. & Verhaar M.A. Effects of cucumber genotype on the induction of resistance to powdery mildew, *Sphaerotheca fuliginea*, by 2,6-dichloroisonicotinic acid, 756
Hock J., Kranz J. & Renfro B.L. Studies on the epidemiology of the tar spot disease complex of maize in Mexico, 490
Hockenhull J. *see* Knudsen I.M.B.
Holmes S.J.I. *see* Dempster L.C.
Hovmøller M.S., Munk L. & stergård H. Comparison of mobile and stationary spore-sampling techniques for estimating virulence frequencies in aerial barley powdery mildew populations, 829

Howell A.B. & Erwin D.C. Characterization and persistence of *Verticillium albo-atrum* isolated from alfalfa growing in high temperature regions of southern California, 734
Hsiang T. *see* Goodwin P.H.
Hughes G. & Madden L.V. Some methods allowing for aggregated patterns of disease incidence in the analysis of data from designed experiments, 927
Hunter P.J. *see* Bevan J.R.
Hussey R.S. *see* Price N.S.

Iacobellis N.S. *see* Caponero A.
Ikegami M. *see* Siriwong P.
Irigoyen J. *see* Aguirreolea J.

Jacks M. *see* Cockerell V.
Jaffe B. *see* Fletcher J.T.
Jamaux I., Gelie B. & Lamarque C. Early stages of infection of rapeseed petals and leaves by *Sclerotinia sclerotiorum* revealed by scanning electron microscopy, 22
Jameson P.E. *see* Eason J.R.
Jansen A.H.J. *see* Smid E.J.
Janssen E.M. *see* Drenth A.
Jeger M.J. *see* Diourte M.
Jenkinson P. *see* Parry D.W.
Jensen D.F. *see* Knudsen I.M.B.
Jensen H.P. *see* Lyngkjør M.F.
Jessop A.C. *see* Brown J.K.M.
Jiménez-D'az R.M. *see* Castillo P., Melero-Vara J.M., Navas-Cortés J.A.
Jones E.E. & Deacon J.W. Mycoparasite-like behaviour of the plant pathogen *Pythium aphanidermatum in vitro*, 396

Kamata N. *see* Sugimura Y.
Kearns L.P. & Hale C.N. Incidence of bacteria inhibitory to *Erwinia amylovora* from blossoms in New Zealand apple orchards, 918
Keck M., Chartier R., Zislavsky W., Lecomte P. & Paulin J.P. Heat treatment of plant propagation material for the control of fire blight, 124
Keel C. *see* Maurhofer M.
Keenan J.P. *see* Walters D.R.
Kemp G.H.J. *see* Viljoen A.
Kennedy D.M. *see* Duncan J.M.
Kenyon D.M., Dixon G.R. & Helfer S. Culture *in vitro* of *Rhododendron* and *Erysiphe* sp., 350
Kittipakorn K. *see* Siriwong P.
Klironomou E. *see* Vakalounakis D.J.
Knudsen I.M.B., Hockenhull J. & Jensen D.F. Biocontrol of seedling diseases of barley and wheat caused by *Fusarium culmorum* and *Bipolaris sorokiniana*: effects of selected fungal antagonists on growth and yield components, 467
Kousik C.S., Snow J.P., Berggren G.T. & Harville B.G. Effect of temperature on virulence of *Rhizoctonia solani* on soybean leaves and seedlings, 580
Kranz J. *see* Hau B., Hock J.
Kuszala C. *see* Seddas A.

Lamarque C. *see* Jamaux I.
Lannou C., de Vallavieille-Pope C. & Goyeau H. Induced resistance in host mixtures and its effect on disease control in computer-simulated epidemics, 478
Larena I. *see* de Cal A.
Leavitt G.M. *see* Duncan R.A.
Lecomte P. *see* Keck M.
Letourmy P. *see* Rott P.
Lewis B.G. *see* Barr K.J.
Liu H.W. *see* Goodwin P.H.
Locke T. & Phillips A.N. The occurrence of carbendazim resistance in *Rhynchosporium secalis* on winter barley in England and Wales in 1992 and 1993, 294
Lowe R. *see* Wallace A.
Lucas J.A. *see* Dyer P.S.
Lundborg T. *see* Olbe M.
Luo Y. & Zeng S.M. Simulation studies on epidemics of wheat stripe rust (*Puccinia striiformis*) on slow-rusting cultivars and analysis of effects of resistance components, 340
Lyngkjør M.F., Jensen H.P. & stergård H. A Japanese powdery mildew isolate with exceptionally large infection efficiency on Mlo-resistant barley, 786
Lyon G.D., Reglinski T. & Newton A.C. Novel disease control compounds: the potential to 'immunize' plants against infection, 407

McCartney H.A. *see* Figueroa L.
MacDonald M.V. *see* Amoah B.K., Day J.P.
Mac'as G. *see* Maddison A.C.
McLeod L. *see* Parry D.W.
McMillan R.T., Jr *see* Ramos L.J.
McPherson A. *see* Walters D.R.
McQuilken M.P., Mitchell S.J., Budge S.P., Whipps J.M. Fenlon J.S. & Archer S.A. Effect of *Coniothyrium minitans* on sclerotial survival and apothecial production of *Sclerotinia sclerotiorum* in field-grown oilseed rape, 883
Madden L.V. *see* Ammar E.D., Hughes G.
Maddison A.C., Mac'as G., Moreira C., Arias R. & Neira R. Cocoa production in Ecuador in relation to dry-season escape from pod rot caused by *Crinipellis perniciosa* and *Moniliophthora roreri*, 982
Mantle P.G. & Swan D.J. Effect of male sterility on ergot disease spread in wheat, 392
Marasas W.F.O. *see* Viljoen A.
Martelli G.P. *see* Boscia D.
Masuko M. *see* Shirane N.
Mäurer R. & Seemüller E. Nature and genetic relatedness of the mycoplasma-like organism causing rubus stunt in Europe, 244
Maurhofer M., Keel C., Haas D. & Défago G. Influence of plant species on disease suppression by *Pseudomonas fluorescens* strain CHA0 with enhanced antibiotic production, 40
Meignoz R. *see* Seddas A.
Melero-Vara J.M., Blanco-Lùpez M.A., Bejarano-Alcázar J. & Jiménez-D'az R.M. Control of *Verticillium* wilt of cotton by means of soil solarization and tolerant cultivars in southern Spain, 250
Melgarejo P. *see* de Cal A.
Mepsted R., Flood J., Paul T., Airede C. & Cooper R.M. A model system for rapid selection for resistance and investigation of resistance mechanisms in *Fusarium* wilt of oil palm, 749
Merdinoglu D. *see* Geyl L.
Migheli Q., Berio T., Gullino M.L. & Garibaldi A. Electrophoretic karyotype variation among pathotypes of *Fusarium oxysporum* f.sp. *dianthi*, 308
Mitchell S.J. *see* McQuilken M.P.
Mol L., Scholte K. & Vos J. Effects of crop rotation and removal of crop debris on the soil population of two isolates of *Verticillium dahliae*, 1070
Moreira C. *see* Maddison A.C.
Morton A., Carder J.H. & Barbara D.J. Sequences of the internal transcribed spacers of the ribosomal RNA genes and relationships between isolates of *Verticillium alboatrum* and *V. dahliae*, 183
Mundt C.C., Brophy L.S. & Schmitt M.S. Disease severity and yield of pure-line wheat cultivars and mixtures in the presence of eyespot, yellow rust, and their combination, 173
Mundt C.C. *see* Ahmed H.U.
Munk L. *see* Hovmøller M.S.
Musabyimana T., Sehene C. & Bandyopadhyay R. Ergot resistance in sorghum in relation to flowering, inoculation technique and disease development, 109
Muthumeenakashi S. *see* Fletcher J.T.

Narayanan K.R. *see* Ramos L.J.
Natsuaki T. *see* Sugimura Y.
Navas-Cortés J.A., Trapero-Casas A. & Jiménez-D'az R.M. Survival of *Didymella rabiei* in chickpea straw debris in Spain, 332
Nayudu M.V. *see* Satyanarayana T.
Neate S.M. *see* Balali G.R.
Neira R. *see* Maddison A.C.
Newton A.C. & Andrivon D. Assumptions and implications of current gene-for-gene hypotheses, 607
Newton A.C. *see* Lyon G.D.
Nicholson P. *see* Amoah B.K.

O'Brien P.A. *see* Yang H.A.
O'Connell M.N. *see* Simpson M.J.A.
Okuda S. *see* Sugimura Y.
Olbe M., Sommarin M., Gustafsson M. & Lundborg T. Effect of the fungal pathogen *Bipolaris sorokiniana* toxin prehelminthosporol on barley root plasma membrane vesicles, 625
Østergård H. *see* Hovmøller M.S., Lyngkjør M.F.

Padayhag B.F. *see* Sugimura Y.
Padgham D.E. *see* Stevenson P.C.
Panagopoulos C.G. *see* Antoniou P.P.
Parker S.R., Shaw M.W. & Royle D.J. The reliability of visual estimates of disease severity on cereal leaves, 856
Parry D.W., Jenkinson P. & McLeod L. *Fusarium* ear blight (scab) in small grain cereals—a review, 207

Pascual S. *see* de Cal A.
Paul T. *see* Mepsted R.
Paulin J.P. *see* Keck M.
Phillips A.N. *see* Locke T.
Prasertkit O. *see* Hau B.
Price N.S., Roncadori R.W. & Hussey R.S. The growth of nematode 'tolerant' and 'intolerant' soyabeans as affected by phosphorus, *Glomus intraradices* and light, 597

Rafin C. & Tirilly Y. Characteristics and pathogenicity of *Pythium* spp. associated with root rot of tomatoes in soilless culture in Brittany, France, 779
Ramage R.A. & Sutherland M.W. High and low pre-inoculation temperatures decrease the effectiveness of the *Lr20* and *Sr15* rust resistance genes in wheat, 772
Ramos L.J., Bharathan N., McMillan R.T., Jr & Narayanan K.R. Histopathological changes associated with silverleaf syndrome in squash, 316
Reardon T.B. *see* Riley I.T.
Reddy D.V.R. *see* Satyanarayana T.
Reddy R.K. *see* Butler D.R.
Reglinski T. *see* Lyon G.D.
Renfro B.L. *see* Hau B., Hock J.
Rennie W.J. *see* Cockerell V.
Resende M.L.V., Flood J. & Cooper R.M. Effect of method of inoculation, inoculum density and seedling age at inoculation on the expression of resistance of cocoa (*Theobroma cacao* L.) to *Verticillium dahliae* Kleb., 374
Reuveni M., Agapov V. & Reuveni R. Suppression of cucumber powdery mildew (*Sphaerotheca fugilinea*) by foliar sprays of phosphate and potassium salts, 31
Reuveni R. *see* Reuveni M.
Rezanoor H.N. *see* Amoah B.K.
Ridout M.S. *see* Xu X.-M.
Riley I.T. & Reardon T.B. Isolation and characterisation of *Clavibacter tritici* associated with *Anguina tritici* in wheat from Western Australia, 805
Roberts J.K. *see* Burdon J.J.
Roderick H.W. & Clifford B.C. Variation in adult plant resistance to powdery mildew in spring oats under field and laboratory conditions, 366
Roncadori R.W. *see* Price N.S.
Rose J. *see* Brasier C.M.
Rosenow D.T. *see* Diourte M.
Rossall S. *see* Fiddaman P.J.
Rotino G.L. *see* Cristinzio G.
Rott P., Soupa D., Brunet Y., Feldmann P. & Letourmy P. Leaf scald (*Xanthomonas albilineans*) incidence and its effect on yield in seven sugarcane cultivars in Guadeloupe, 1075
Royle D.J. *see* Parker S.R.

Sache I. & Zadoks J.C. Effect of rust (*Uromyces viciae-fabae*) on yield components of faba bean, 675
Salaverri J. *see* Aguirreolea J.
Sánchez-D'az M. *see* Aguirreolea J.
Sanoamuang N. & Gaunt R.E. Persistence and fitness of carbendazim- and dicarboximide-resistant isolates of *Monilinia fructicola* (Wint.) Honey in flowers, shoots and fruit of stone fruit, 448
Satyanarayana T., Broglio E.P., Sreenivasulu P., Reddy D.V.R. & Nayudu M.V. Genome characterization of a new strain of peanut chlorotic streak virus causing chlorotic vein banding disease of groundnut (*Arachis hypogaea* L.) in India, 966
Savino V. *see* Boscia D.
Scala F. *see* Castoria R., Cristinzio G.
Schmitt M.S. *see* Mundt C.C.
Scholte K. *see* Mol L.
Scott E.S. *see* Balali G.R.
Seddas A., Meignoz R., Kuszala C. & Boudon-Padieu E. Evidence for the physical integrity of flavescence dorée phytoplasmas purified by immunoaffinity from infected plants or leafhoppers and the plant pathogenicity of phytoplasmas from leafhoppers, 971
Seemüller E. *see* Mäurer R.
Sehene C. *see* Musabyimana T.
Shattock R.C. *see* Al-Kherb S.M.
Shaw D.S. *see* Al-Kherb S.M.
Shaw M.W. *see* Figueroa L., Parker S.R.
Shirane N., Masuko M. & Takeda R. Effects of SSF-126, a novel alkoxyiminoacetamide blasticide, on mycelial growth and oxygen consumption of *Pyricularia oryzae*, 636
Simpson M.J.A., O'Connell M.N., Stobart R.M. & Cullum F.J. Occurrence and viability of *Septoria oenotherae* in commercial seed samples of evening primrose, 547

Siriwong P., Kittipakorn K. & Ikegami M. Characterization of chilli vein-banding mottle virus isolated from pepper in Thailand, 718
Sivasithamparam K. *see* Willamson P.M., Yang H.A.
Smid E.J., Jansen A.H.J. & Gorris L.G.M. Detection of *Erwinia carotovora* subsp. *atroseptica* and *Erwinia chrysanthemi* in potato tubers using polymerase chain reaction, 1058
Snow J.P. *see* Kousik C.S.
Sommarin M. *see* Olbe M.
Soupa D. *see* Rott P.
Spence N.J. & Walkey D.G.A. Variation for pathogenicity among isolates of bean common mosaic virus in Africa and a reinterpretation of the genetic relationship between cultivars of *Phaseolus vulgaris* and pathotypes of BCMV, 527
Spotts R.A. & Cervantes L.A. Factors affecting the severity of bacterial canker of pear caused by *Pseudomonas syringae* pv. *syringae*, 325
Sreenivasulu P. *see* Satyanarayana T.
Stack J.P. *see* Diourte M.
Stapleton J.J. *see* Duncan R.A.
Starr J.L. *see* Diourte M.
Stenlid J. *see* Swedjemark G.
Stevenson P.C., Padgham D.E. & Haware M.P. Root exudates associated with the resistance of four chickpea cultivars (*Cicer arietinum*) to two races of *Fusarium oxysporum* f.sp. *ciceri*, 686
Stobart R.M. *see* Simpson M.J.A.
Strange R.N. *see* Edwards C.
Sugimura Y., Padayhag B.F., Ceniza M.S., Kamata N., Eguchi S., Natsuaki T. & Okuda S. Essential oil production increased by using virus-free patchouli plants derived from meristem-tip culture, 510
Sun Zhi-Huan *see* Dai Qun
Sutherland M.W. *see* Ramage R.A.
Swan D.J. *see* Mantle P.G.
Swedjemark G. & Stenlid J. Susceptibility of conifer and broadleaf seedlings to Swedish S and P strains of *Heterobasidion annosum*, 73

Takeda R. *see* Shirane N.
Taylor J.D. *see* Bevan J.R.
Than K.A. *see* Willamson P.M.
Tian S.P. & Bertolini P. Effects of low temperature on mycelial growth and spore germination of *Botrytis allii* in culture and on its pathogenicity to stored garlic bulbs, 1008
Tirilly Y. *see* Rafin C.
Tjamos E.C. *see* Antoniou P.P., Elena K.
Trapero-Casas A. *see* Navas-Cortés J.A.

Vakalounakis D.J. Inheritance and linkage of resistance in cucumber line SMR-18 to races 1 and 2 of *Fusarium oxysporum* F.sp. *cucumerinum*, 169
Vakalounakis D.J. & Klironomou E. Race and mating type identification of powdery mildew on cucurbits in Greece, 1033
Valentin P. *see* Geyl L.
Valverde R.A. *see* Arancibia R.A.
van den Bosch F. *see* Habtu Assefa
van Santen G. *see* Xu X.-M.
Verhaar M.A..*see* Hijwegen T.
Viljoen A., Wingfield M.J., Kemp G.H.J. & Marasas W.F.O. Susceptibility of pines in South Africa to the pitch canker fungus *Fusarium subglutinans* f.sp. *pini*, 877
Virányi F. & Gulya T.J. Inter-isolate variation for virulence in *Plasmopara halstedii* (sunflower downy mildew) from Hungary, 619
Vivian A. *see* Bevan J.R.
Vos J. *see* Mol L.
Vovlas N. *see* Castillo P.

Wadia K.D.R. *see* Butler D.R.
Walkey D.G.A. *see* Spence N.J.
Wallace A., Williams N.A., Lowe R. & Harrison J.G. Detection of *Spongospora subterranea* using monoclonal antibodies in ELISA, 355
Walsh M. *see* Cassells A.C.
Walters D.R., Keenan J.P., Cowley T., McPherson A. & Havis N.D. Inhibition of polyamine biosynthesis in *Phytophthora infestans* and *Pythium ultimum*, 80
Welham S.J. *see* Figueroa L.
Whipps J.M. *see* McQuilken M.P.
Whisson D.L. *see* Balali G.R.

Whitehead A.G. Decline of potato cyst nematodes, *Globodera rostochiensis* and *G. pallida*, in spring barley microplots, 191
Wicks T.J. *see* Balali G.R.
Williams N.A. *see* Wallace A.
Willamson P.M., Than K.A., Sivasithamparam K., Cowling W.A. & Edgar J.A. Detection of resistance to *Diaporthe toxica* in asymptomatically infected lupin seedlings based on an immunoassay for phomopsin, 95
Wingfield M.J. *see* Viljoen A.
Wisniewski M., Droby S., Chalutz E. & Eilam Y. Effects of Ca^{2+} and Mg^{2+} on *Botrytis cinerea* and *Penicillium expansum in vitro* and on the biocontrol activity of *Candida oleophila*, 1016
Wolfe M.S. New honorary member of the British Society for Plant Pathology, 763
Wright D.M. *see* Fletcher J.T.

Xu X.-M., Butt D.J. & Ridout M.S. Temporal patterns of airborne conidia of *Podosphaera leucotricha*, causal agent of apple powdery mildew, 944
Xu X.-M., Butt D.J. & van Santen G. A dynamic model simulating infection of apple leaves by *Venturia inaequalis*, 865
Xue B.G. *see* Goodwin P.H.

Yang H.A., Sivasithamparam K., Barton J.E. & O'Brien P.A. Characterization of cereal bare patch isolates of *Rhizoctonia solani* by random amplified polymorphic DNA analysis, 811

Zadoks J.C. *see* Habtu Assefa, Sache I.
Zeng S.M. *see* Luo Y.
Zislavsky W. *see* Keck M.
Zoina A. *see* Castoria R.